

## Role of *dprA* in transformation of *Campylobacter jejuni*

Tohru Takata<sup>a,\*</sup>, Takafumi Ando<sup>a,1</sup>, Dawn A. Israel<sup>c</sup>,  
Trudy M. Wassenaar<sup>d</sup>, Martin J. Blaser<sup>a,b</sup>

<sup>a</sup> Department of Medicine, New York University School of Medicine, New York, NY, USA

<sup>b</sup> Department of Microbiology, New York University School of Medicine, New York, NY, USA

<sup>c</sup> Division of Gastroenterology, Department of Medicine, Vanderbilt University School of Medicine, Nashville, TN, USA

<sup>d</sup> Molecular Microbiology and Genomics Consultants, Zotzenheim, Germany

Received 10 May 2005; received in revised form 27 August 2005; accepted 29 August 2005

First published online 19 September 2005

Edited by A.H.M. van Vliet

### Abstract

The role of a *dprA* ortholog (Cj0634) in *Campylobacter jejuni* transformation was phenotypically assessed using two strains. *C. jejuni* strain 11168 was naturally competent for transformation by chromosomal DNA, while efficiency decreased 100-fold in a Cj0634::*aphA* mutant, whereas *C. jejuni* strain 480 was not naturally competent. *C. jejuni* strain 480 but not 11168 could be electro-transformed by shuttle plasmid pRY111, an effect completely abolished by Cj0634 interruption. Complementation of the Cj0634 mutation in *C. jejuni* strain 480 *in trans* with vectors containing the *dprA* homologs from *C. jejuni*, *Helicobacter pylori*, or *Haemophilus influenzae*, completely (for Cj0634) or partially (*H. pylori* > *H. influenzae*) restored electro-transformation. Thus, *C. jejuni* expresses a DprA ortholog that functionally most closely resembles that of *H. pylori* and is involved in DNA transformation.

© 2005 Federation of European Microbiological Societies. Published by Elsevier B.V. All rights reserved.

**Keywords:** Transformation; *dprA*; Competence; *Campylobacter jejuni*

### 1. Introduction

*Campylobacter jejuni* is a gram negative microaerophilic bacterium that colonizes the gastrointestinal tract in avian and mammalian species, and is a common cause of diarrheal diseases in humans [1]. The genetically heterogeneous *C. jejuni* are naturally competent to take up and integrate DNA [2,3]; this heterogeneity may result, at least in part, from intraspecies horizontal gene transfer. The ability for horizontal DNA transfer may be an

important characteristic enabling survival of *C. jejuni* in various natural hosts. Characterization of the factors involved in *C. jejuni* DNA transfer may lead to insights that are generally applicable to other species.

Prior work by our group had drawn our attention to DprA as one of the factors involved in bacterial uptake and integration of DNA. Analysis of the complete genome sequence of *C. jejuni* 11168 [4] revealed a 774-bp open reading frame (Cj0634) predicted to encode a protein with homology to DprA (encoded by *dprA*) of *Haemophilus influenzae* and *Helicobacter pylori*. In *H. influenzae*, DprA is essential for transformation by chromosomal DNA [5], whereas in *H. pylori*, DprA facilitates but is not essential for transformation by chromosomal DNA, and is necessary for plasmid uptake [6,7].

\* Corresponding author. Present address: First Department of Internal Medicine, School of Medicine, Fukuoka University, Fukuoka 814-0180, Japan. Tel.: +81 92 801 1011; fax: +81 92 865 5656.

E-mail address: takattol@cis.fukuoka-u.ac.jp (T. Takata).

<sup>1</sup> Present address: Department of Therapeutic Medicine, Nagoya University Graduate School of Medicine, Nagoya, Japan.